

Help find relief from your severe and persistent pain*

Your healthcare provider chose BELBUCA, a long-acting opioid, for when other options are inadequate

Potential savings information on last page.

*APPROVED USE

BELBUCA® (buprenorphine buccal film) CIII is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage severe and persistent pain that requires an extended treatment period with a daily opioid pain medicine when other pain medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not to be taken on an “as needed” basis.

IMPORTANT SAFETY INFORMATION about BELBUCA®

WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF BELBUCA

Addiction, Abuse, and Misuse

BELBUCA exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing BELBUCA and monitor regularly for these behaviors and conditions.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of BELBUCA, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of BELBUCA are essential. Misuse or abuse of BELBUCA by chewing, swallowing, snorting, or injecting buprenorphine extracted from the buccal film will result in the uncontrolled delivery of buprenorphine and poses a significant risk of overdose and death.

Accidental Exposure

Accidental exposure to even one dose of BELBUCA, especially in children, can result in a fatal overdose of buprenorphine.

Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of BELBUCA and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.

Neonatal Opioid Withdrawal Syndrome (NOWS)

Advise pregnant women using opioids for an extended period of time of the risk of Neonatal Opioid Withdrawal Syndrome, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription

Please see full [Prescribing Information](#), including Boxed Warning on Addiction, Abuse, and Misuse, and other serious risks, and Medication Guide.

Speak to your healthcare provider if you have questions about BELBUCA.

Why has BELBUCA been prescribed for me?

Your healthcare provider is starting you on BELBUCA, a long-acting opioid, because your pain isn't well managed



Long-acting opioids may be used when:



Your pain causes nighttime awakenings




Your current treatment requires frequent daily doses of medicine



Your pain requires around-the-clock relief



Your current treatment causes tolerability issues and/or side effects



BELBUCA is a strong prescription pain medicine (buprenorphine)

BELBUCA contains an opioid (narcotic) that is used to manage severe and persistent pain that requires an extended treatment period with a daily opioid pain medicine. BELBUCA is used when other pain medicines do not treat your pain well enough or you cannot tolerate them.

IMPORTANT SAFETY INFORMATION about BELBUCA® (cont'd)

Important information about BELBUCA:

- **Get emergency help or call 911 right away if you take too much BELBUCA (overdose).** When you first start taking BELBUCA, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Ask your healthcare provider about medicines like naloxone or nalmefene that can be used in an emergency to reverse an opioid overdose.
- Taking BELBUCA with other opioid medicines, benzodiazepines, gabapentinoids (gabapentin or pregabalin), alcohol, or other central nervous system

depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.

- Never give anyone else your BELBUCA. They could die from taking it. Selling or giving away BELBUCA is against the law.
- Store BELBUCA securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

Please see full [Prescribing Information](#), including **Boxed Warning on Addiction, Abuse, and Misuse**, and other serious risks, and **Medication Guide**.
Speak to your healthcare provider if you have questions about BELBUCA.

How is BELBUCA different?

BELBUCA is the first and only schedule III opioid using buprenorphine buccal film technology



BELBUCA is a long-acting opioid

BELBUCA works over an extended period of time, which means it stays in your system longer than short-acting opioids. BELBUCA will be taken every 12 hours, instead of every 4 to 6 hours like many short-acting opioids.



For illustrative purposes only.
Not the actual product or size.

BELBUCA uses a different delivery system

BELBUCA is a thin, square film that you put in your mouth and press against the inside of your cheek. The film delivers the medicine, buprenorphine, efficiently through the inside of your cheek, entering your bloodstream more directly.

IMPORTANT SAFETY INFORMATION about BELBUCA® (cont'd)

Do not use BELBUCA if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

Before taking BELBUCA, tell your healthcare provider if you have a history of:

- head injury, seizures
- heart rhythm problems (long QT syndrome)
- liver, kidney, thyroid problems
- pancreas or gallbladder problems
- problems urinating
- tooth problems, including a history of cavities
- abuse of street or prescription drugs, alcohol addiction, opioid overdose, or mental health problems

Please see full [Prescribing Information](#), including **Boxed Warning on Addiction, Abuse, and Misuse**, and other serious risks, and **Medication Guide**.
Speak to your healthcare provider if you have questions about BELBUCA.

What can I expect from BELBUCA?



BELBUCA has been proven to reduce severe and persistent pain*

- In people previously taking commonly prescribed opioids



The possible side effects of BELBUCA include nausea, constipation, headache, vomiting, dizziness, and sleepiness*

- Call your healthcare provider if you have any of these symptoms and they are severe



BELBUCA has 7 dosage strengths and is taken 2 times a day


- Multiple dosage strengths will help you and your healthcare provider find the dose that provides the severe and persistent pain relief you need

As you begin treatment

It is important to know that your starting dose may not be the final dose

It may take time for you and your doctor to find the optimal dose for you.

BELBUCA offers 7 dosage strengths



*Results from clinical studies compared with placebo (sugar pill). Individual results may vary.

IMPORTANT SAFETY INFORMATION about BELBUCA® (cont'd)

Tell your healthcare provider if you are:

- **noticing your pain getting worse.** If your pain gets worse after you take BELBUCA, do not take more of BELBUCA without first talking to your healthcare provider. Talk to your healthcare provider if the pain that you have increases, if you feel more sensitive to pain, or if you have new pain after taking BELBUCA.
- **pregnant or planning to become pregnant.** Prolonged use of BELBUCA during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** Not recommended during treatment with BELBUCA. It may harm your baby.
- living in a household where there are small children or someone who has abused street or prescription drugs.

- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking BELBUCA with certain other medicines can cause serious side effects and could lead to death.

When taking BELBUCA:

- Do not change your dose. Apply BELBUCA exactly as prescribed by your healthcare provider. Use the lowest effective dose possible for the shortest time needed.
- See the detailed Instructions for Use for information about how to apply BELBUCA.
- Do not apply BELBUCA if the package seal is broken or the film is cut, damaged, or changed in any way.

Please see full [Prescribing Information](#), including [Boxed Warning on Addiction, Abuse, and Misuse](#), and other serious risks, and [Medication Guide](#).
Speak to your healthcare provider if you have questions about BELBUCA.

Taking BELBUCA

Make sure you apply the film at the same times each day, following your healthcare provider's instructions

Before you begin, wet the inside of your cheek with your tongue or with water.



1. PEEL OPEN PACKAGE

- With clean, dry fingers, peel open the foil package. Fold along the dotted line at the top of the package, and tear at the perforation
- You can also use scissors to carefully cut along the dotted line



2. PLACE ON FINGER

- Carefully remove the BELBUCA film from the foil package
- Place the film on your dry finger with the yellow side facing up



3. PRESS INSIDE OF CHEEK

- After wetting the inside of your cheek with your tongue or by rinsing with water, press the yellow side against the inside of your moistened cheek. Hold it in place for 5 seconds, and then remove your finger
- Leave BELBUCA on the inside of your cheek until fully dissolved, usually within 30 minutes
- After BELBUCA is completely dissolved, rinse your mouth with water, and swallow. Wait at least one hour before brushing your teeth

- Avoid eating food or drinking liquids until BELBUCA has dissolved

- Avoid touching or moving BELBUCA with your tongue or finger after it is in place

- Do not chew or swallow BELBUCA

LEARN MORE ABOUT APPLYING BELBUCA

by viewing the **application video** at belbuca.com/resources

IMPORTANT SAFETY INFORMATION about BELBUCA® (cont'd)

When taking BELBUCA: (cont'd)

- After the film has adhered to your cheek, avoid eating or drinking until the film has completely dissolved, usually within 30 minutes.
- After BELBUCA is completely dissolved, rinse your mouth with water and swallow. Wait for at least one hour before brushing teeth.

Please see full [Prescribing Information](#), including **Boxed Warning on Addiction, Abuse, and Misuse**, and other serious risks, and **Medication Guide**. Speak to your healthcare provider if you have questions about BELBUCA.

IMPORTANT SAFETY INFORMATION about BELBUCA®

Important information about BELBUCA:

- **Get emergency help or call 911 right away if you take too much BELBUCA (overdose).** When you first start taking BELBUCA, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Ask your healthcare provider about medicines like naloxone or nalmefene that can be used in an emergency to reverse an opioid overdose.
- Taking BELBUCA with other opioid medicines, benzodiazepines, gabapentinoids (gabapentin or pregabalin), alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your BELBUCA. They could die from taking it. Selling or giving away BELBUCA is against the law.
- Store BELBUCA securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

Do not use BELBUCA if you have:

- severe asthma, trouble breathing, or other lung problems
- a bowel blockage or have narrowing of the stomach or intestines.

Before taking BELBUCA, tell your healthcare provider if you have a history of:

- head injury, seizures
- heart rhythm problems (long QT syndrome)
- liver, kidney, thyroid problems
- pancreas or gallbladder problems
- problems urinating
- tooth problems, including a history of cavities
- abuse of street or prescription drugs, alcohol addiction, opioid overdose, or mental health problems

Tell your healthcare provider if you are:

- **noticing your pain getting worse.** If your pain gets worse after you take BELBUCA, do not take more of BELBUCA without first talking to your healthcare provider. Talk to your healthcare provider if the pain that you have increases, if you feel more sensitive to pain, or if you have new pain after taking BELBUCA.
- **pregnant or planning to become pregnant.** Prolonged use of BELBUCA during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** Not recommended during treatment with BELBUCA. It may harm your baby.
- living in a household where there are small children or someone who has abused street or prescription drugs.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking BELBUCA with certain other medicines can cause serious side effects and could lead to death.

When taking BELBUCA:

- Do not change your dose. Apply BELBUCA exactly as prescribed by your healthcare provider. Use the lowest effective dose possible for the shortest time needed.
- See the detailed Instructions for Use for information about how to apply BELBUCA.

- Do not apply BELBUCA if the package seal is broken or the film is cut, damaged, or changed in any way.
- After the film has adhered to your cheek, avoid eating or drinking until the film has completely dissolved, usually within 30 minutes.
- After BELBUCA is completely dissolved, rinse your mouth with water and swallow. Wait for at least one hour before brushing teeth.
- Report any problems with your teeth immediately to your healthcare provider and schedule an appointment with a dentist. Tell your dentist that you have started taking BELBUCA.
- Avoid touching or moving the buccal film with your tongue or fingers.
- **Do not chew, swallow, snort or inject BELBUCA. This will result in uncontrolled delivery of buprenorphine and may cause you to overdose and die.**
- **Call your healthcare provider if the dose you are using does not control your pain.**
- **Do not stop using BELBUCA without talking to your healthcare provider.**
- Dispose of expired, unwanted, or unused BELBUCA by removing the BELBUCA film from the foil packaging, and promptly flushing down the toilet (if a drug takeback option is not readily available). Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines.

While using BELBUCA DO NOT:

- Drive or operate heavy machinery, until you know how BELBUCA affects you. BELBUCA can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines containing alcohol. Using products containing alcohol during treatment with BELBUCA may cause you to overdose and die.

The possible side effects of BELBUCA are:

- nausea, constipation, headache, vomiting, dizziness, and sleepiness. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help or call 911 right away if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of BELBUCA. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. For more information go to dailymed.nlm.nih.gov.

Please see full [Prescribing Information](#), including [Boxed Warning on Addiction, Abuse, and Misuse](#), and other serious risks, and [Medication Guide](#).

Pay as little as \$0 with the **BELBUCA** savings card

Eligible, commercially-insured
patients may start saving
right away



*Eligibility Criteria

This offer is valid for the purchase of BELBUCA manufactured for Collegium Pharmaceutical, Inc. and lawfully purchased from an authorized retailer or distributor in the United States or its territories. **This offer is not insurance and is not valid for mail order or prescriptions purchased under Medicaid, Medicare, TRICARE, or similar federal or state programs, or for patients who are Medicare eligible and enrolled in an employer-sponsored group waiver health plan or government-subsidized prescription drug benefit program for retirees.** Offer not valid where prohibited by law, taxed, or restricted. Your offer has its own unique ID number. This offer is not transferable, is limited to one per person, and may not be combined with any other offer. Offer must be presented along with a valid prescription for BELBUCA at the time of purchase. If you lose your card before the first time you use it, simply print or download a new copy at [BELBUCA.com](https://www.belbucacard.com) and bring it to the pharmacy. Cash Discount Cards and other non-insurance plans are not valid as primary under this offer. If the patient is eligible for drug benefits under any such program, the patient cannot use this offer. By using this offer, the patient certifies that he or she will comply with any terms of his or her health insurance contract requiring notification to his or her payer of the existence and/or value of this offer. This offer is not valid for cash paying patients and/or patients under 18 years of age.

Collegium Pharmaceutical, Inc. reserves the right to change or discontinue this offer at any time without notice.

[EXPLORE POTENTIAL SAVINGS AT BELBUCA.COM/SAVINGS](https://www.belbucacard.com/savings)

Please see full [Prescribing Information](#), including **Boxed Warning on Addiction, Abuse, and Misuse, and other serious risks, and Medication Guide**.
Speak to your healthcare provider if you have questions about BELBUCA.

Rx Only

BELBUCA is a registered trademark of BioDelivery Sciences International, Inc., a wholly owned subsidiary of Collegium Pharmaceutical, Inc.
© 2026 Collegium Pharmaceutical, Inc. All rights reserved.
PP-BEL-US-0211 05/26